

RESEARCH ARTICLE

Open Access

Osteogenic differentiation of amniotic epithelial cells: synergism of pulsed electromagnetic field and biochemical stimuli

Qian Wang^{1,2,3,4}, Wenchao Wu³, Xiaoyu Han³, Ai Zheng⁵, Song Lei⁶, Jiang Wu⁷, Huaiqing Chen^{3,7}, Chengqi He^{1,2,4}, Fengming Luo^{3,8} and Xiaojing Liu^{3,1,2*}

Abstract

Background: Pulsed electromagnetic field (PEMF) is a non-invasive physical therapy used in the treatment of fracture nonunion or delayed healing. PEMF can facilitate the osteogenic differentiation of bone marrow mesenchymal stem cells *in vitro*. Amniotic epithelial cells (AECs) have been proposed as a potential source of stem cells for cell therapy. However, whether PEMF could modulate the osteogenic differentiation of AECs is unknown. In the present study, the effects of PEMF on the osteogenic differentiation of AECs were investigated.

Methods: AECs were isolated from amniotic membrane of human placenta by trypsin digestion and were induced by PEMF and/or osteo-induction medium. After 21 days we used real time RT-PCR and immunocytochemistry to study the expression of osteoblast markers. The signal transduction of osteogenesis was further investigated.

Results: The PEMF stimulation, or osteo-induction medium alone could induce osteogenic differentiation of AECs, as shown by expression of osteoblast specific genes and proteins including alkaline phosphatase and osteocalcin. Furthermore, a combination of PEMF and osteo-induction medium had synergy effects on osteogenic differentiation. In our study, the gene expression of BMP-2, Runx2, β -catenin, Nrf2, Keap1 and integrin β 1 were up-regulated in the osteogenic differentiation of AECs induced by PEMF and/or osteo-induction medium.

Conclusions: Combined application of PEMF and osteo-induction medium is synergistic for the osteogenic differentiation of AECs. It might be a novel approach in the bone regenerative medicine.

Keywords: Amniotic epithelial cells, Osteogenic differentiation, Pulsed electromagnetic field, BMP-2, Wnt/ β -catenin signaling, Reactive oxygen species, Integrin β 1

Background

Amniotic epithelial cells (AECs), derived from the placenta, possess several advantages over both embryonic stem cells (ESCs) and adult stem cells. They express ESCs markers such as SSEA-1, SSEA-4 and Oct-4, and have the ability to differentiate into all three germ layers *in vitro* [1,2]. Therefore, AECs have been proposed to be a good candidate for cell transplantation and regenerative medicine [3,4].

AECs can be induced to differentiate into osteoblasts *in vitro* by treating with osteo-induction medium, which contains biochemical factors including dexamethasone, β -glycerol phosphate and ascorbic acid [5]. However, efficient induction of AECs differentiation into osteoblasts remains a challenge, and its mechanism is not fully understood.

Pulsed electromagnetic field (PEMF), a non-invasive physical treatment, is now used clinically to promote bone healing for fracture nonunion or delayed fracture healing [6,7]. PEMF has various biological functions and can affect bone metabolism. Several studies have shown that PEMF can facilitate the osteogenesis by its direct effects on osteoblasts [8,9]. Recently, it has been reported that PEMF with specific parameters could modulate

* Correspondence: xiaojingliu67@gmail.com

³Laboratory of Cardiovascular Diseases, Regenerative Medicine Research Center, West China Hospital, Sichuan University, Chengdu, Sichuan, P. R. China

¹Center of Rehabilitation Medicine, West China Hospital, Sichuan University, Chengdu, Sichuan, P. R. China

²Rehabilitation Key Laboratory of Sichuan Province, Chengdu, Sichuan, P. R. China
Full list of author information is available at the end of the article

osteogenic differentiation of bone marrow derived mesenchymal stem cells (BMSCs) *in vitro* [10-12]. These findings suggest that PEMF might be able to induce AECs to differentiate into osteoblasts.

In this study, we tested the hypothesis that PEMF could modulate the osteogenic differentiation of AECs. We hypothesized that physical force (PEMF) and biochemical treatment and/or their combination might play important roles in the process of osteogenic differentiation of AECs. We explored the possible mechanisms, especially the regulating role of BMP-2 and ROS pathways in the process.

Methods

AECs isolation, culture and identification

Five placenta samples through cesarean delivery were collected aseptically with the informed consents of parturients. Primary cultures of human AECs were isolated from amniotic membrane by trypsin digestion method and cultured in standard culture medium, according to Miki's methods [1]. AECs were identified by the specific epithelial cell marker cytokerin 19 using immunocytochemistry. Phenotype of human AECs was analyzed by flow cytometry.

The study was conducted according to the Declaration of Helsinki and approved by the medical ethics committee of the West China Hospital, Sichuan University.

PEMF stimulation

PEMF was generated by a commercial, clinically approved PEMF system (Model XT-2000B). During each pulse, the applied field increased from 0 to 1mT in 1.5 ms and then decayed back to 0 in 5 ms. The 50Hz repetitive pulsed waveform was based on other previous investigations [13,14]. AECs from the third to fifth passages were maintained in standard culture medium as controls or in osteo-induction medium (OM) [5]. For PEMF treatment, the cells were exposed daily to 50Hz 1mT PEMF stimulation for 30 minutes each time and twice a day, with an interval of 12 hours. The treatment lasted for 21 days. Each experiment was in three replicates.

Detection of alkaline phosphatase (ALP) activity by histochemistry staining

The activity of ALP was detected by histochemistry staining using a BCIP/NBT ALP kit (R&D Biotech, UK) according to manufacturer's instructions [15]. The percentage of ALP-positive cells was calculated by ImageJ software (National Institutes of Health, USA).

Detection of osteocalcin (OC) protein expression by immunocytochemistry

AECs were plated onto coverslips in 6-well plates and treated with different stimuli. The protein expression of

OC was measured by immunocytochemistry as described before [16]. Images were captured on a Leica DFC 300FX Digital Camera system (Leica, UK) and analyzed with the ImageJ software to semi-quantitatively determine the level of cytoplasmic OC.

Assessment of calcium deposition by alizarin red staining

Calcium deposition, one of the markers of osteogenic differentiation of stem cell, was assessed by alizarin red S staining [17]. Images were captured on a Leica DFC 300FX Digital Camera system (Leica, UK). The amount of calcified deposition was semi-quantitatively calculated with ImageJ software.

Quantitative reverse transcription PCR (Real-time RT-PCR)

Total RNA was extracted from AECs with TRIZOL (Invitrogen, USA), and cDNA was synthesized using a reverse transcription (RT) kit (Toyobo, Osaka, Japan) [16]. Quantitative PCR was carried out on BIO-RAD CFX96™ Real-Time PCR Detection System with fluorescence dye EvaGreen (EvaGreen Supermix kit, Bio-Rad, USA). Primer sequences are shown in Table 1. Data analysis was carried out by Bio-Rad software using relative quantification. The comparative cycle-threshold method ($2^{-\Delta\Delta CT}$) was used for quantification of gene expression [17]. For quantification, the target sequence was normalized to the β -actin mRNA levels.

Statistical analysis

The experimental data were presented as means \pm SD. Group means were compared by *One-way ANOVA* using the statistical software SPSS 10.0 for Windows (Chicago, IL, USA), and *P* value < 0.05 was considered to be statistically significant.

Results

Characterization and phenotype of isolated AECs

Amniotic epithelial cells (AECs) were successfully isolated from human placenta and formed a confluent monolayer of cobblestone-shaped epithelial cells after 3 days of culturing in the standard culture media (Figure 1A).

In our study, flow cytometry analysis revealed that 54.2% of primary cultured AECs were positive for SSEA-4 and 92.8% were positive for Oct-4 (Figure 1B). Furthermore, the AECs could also express cytokeratin 19, the epithelial cell marker (Figure 1C). Consistent with previous reports [2,5], we provided further evidence that AECs expressed not only the epithelial cell marker cytokeratin 19, but also the stem cell markers such as SSEA-4 and Oct-4.

PEMF modulates the osteogenic differentiation of AECs

In order to examine whether PEMF could play a role in osteogenic differentiation of AECs, we used real time

Table 1 Primer sequences and amplicon sizes for real-time RT-PCR

Genes	Gene Bank ID	Primer Sequence (5'-3')	Amplicon (bp)
ALP	NM_000478	F: 5' GGAACTCCTGACCCTT GACC 3'	86
		R: 5' TCCTGTTTCAGCTCGTA CTGC 3'	
OC	NM_001199662	F: 5' AGGGCAGCGAGGTAG TGAA 3'	151
		R: 5' TCCTGAAAGCCGATGTG GT 3'	
BMP-2	NM_001200	F: 5' TCAAGCCAAACACAAA CAGC 3'	103
		R: 5' AGCCACAATCCAGTC ATTCC 3'	
RUNX2	NM_001015051	F: 5' TTACTTACACCCCGCC AGTC 3'	139
		R: 5' TATGGAGTGCTGCTGG TCTG 3'	
β -catenin	NM_020248	F: 5' CCTATGCAGGGGTGGT CAAC 3'	95
		R: 5' CGACCTGGAAAACGC CATCA 3'	
Integrin β 1	NM_002211	F: 5' CAAGCAGGGCCAAATT GTGG 3'	121
		R: 5' TGTCATCTGGAGGGC AACCC 3'	
Nrf2	NM_001145413	F: 5' ATTGCCTGTAAGTCCT GGTCA 3'	182
		R: 5' ACTGCTCTTTGGACA TCATTTCG 3'	
Keap1	NM_203500	F: 5' GTGGCTGCTCTCAATC GTCT 3'	127
		R: 5' GGATGGTGTTCATT GCTGTG 3'	
β -actin	NM_001099771	F: 5' ACTATCGGCAATGAG CGGTC 3'	77
		R: 5' ATGCCACAGGATTC CATACCC 3'	

RT-PCR and immunocytochemistry to study the expression of osteoblast markers, such as alkaline phosphatase (ALP) and osteocalcin (OC) in AECs followed by (1) standard culture media (control group), (2) administration of PEMF for AECs cultured in standard culture media (PEMF only), (3) osteo-induction medium (OM only), (4) combined treatments with PEMF and osteo-induction medium for various time points (up to 21 days).

Real time RT-PCR results showed that application of PEMF alone increased the gene expression of ALP and OC above basal levels after 7 days of treatment. The osteo-induction medium alone induced the gene expression of ALP and OC at day 3 post-treatment. The combined induction of ALP and OC mRNA expression were

both peaked at day 11, and the elevated level lasted up to day 21 after the combined treatments (Figure 2).

Next, we examined the administration of PEMF and/or OM-induced change of ALP and OC protein level in AECs. Exposure of AECs to PEMF alone induced the ALP activity at day 7. Culture of AECs in osteo-induction medium increased the ALP activity, which reached the highest level at day 11, and the increased level lasted up to day 21 post-treatment. Moreover, combined application of PEMF and osteo-induction medium resulted in a much higher level of ALP activity compared with other groups (Figure 3). The expression of OC protein was analyzed by immunocytochemistry, which showed that AECs after combined stimulation of PEMF and osteo-induction medium presented a much stronger OC protein expression, than what either treatment alone (Figure 4).

The osteogenic differentiation of AECs was also validated by detection of the calcium deposits [17]. Alizarin red S staining results showed that, application of PEMF alone increased calcium deposits in AECs as early as day 7 after treatment. The osteo-induction medium also evoked a marked increase of extracellular matrix calcification in AECs. Furthermore, a combination of PEMF with osteo-induction medium induced much more calcium deposition in AECs than the other treatments, and the effect was peaked at day 11 (Figure 5).

Taken together, we demonstrated that a combination of PEMF and osteo-induction medium had much stronger effects on osteogenic differentiation of AECs, than what either treatment alone had.

PEMF might modulate the osteogenic differentiation of AECs via BMP-2/Runx2 or Wnt/ β -catenin signaling

To investigate the possible role of BMP-2/Runx2 or Wnt/ β -catenin signaling in osteoblast differentiation of AECs, we assessed the expression change of BMP-2, Runx2 and β -catenin mRNA in AECs with different treatments. Exposure of PEMF alone induced the expression of BMP-2, Runx2 and β -catenin above basal levels after 7 days of treatment. Osteo-induction medium alone promoted the expression of BMP-2, Runx2 and β -catenin, followed by a gradual decrease after treatment. Moreover, combined application of PEMF with osteo-induction medium led to a significant up-regulation in the expression of BMP-2, Runx2 and β -catenin compared with other treatments (Figure 6A-C).

Nrf2 and Keap1 might be involved in the PEMF-induced osteogenic differentiation of AECs

Reactive oxygen species (ROS) could play vital role in the self-renewal and differentiation of stem cells [18,19]. To test whether Nrf2 and Keap1, the master regulators of ROS generation, might be implicated in the osteogenic differentiation of AECs, we examined the gene




Figure 1 Characterization and phenotype of isolated Amniotic epithelial cells (AECs). (A) Primary culture of human AECs. AECs were successfully isolated from human term placenta and formed a confluent monolayer of cobblestone-shaped epithelial cells after 3 days of culture in the standard culture media (magnification of 200x). (B) Surface antigen SSEA-4 and the pluripotency marker Oct-4 in primary cultured AECs. Flow cytometry revealed the presence of SSEA-4 (54.2%) and Oct-4 (92.8%) in primary cultured AECs. The open peaks show the isotype-matched antibody control. (C) Specific marker of epithelial cells identified by immunocytochemistry. Primary cultured AECs could express cyokeratin 19, the epithelial cell marker (magnification of 200x).




Figure 2 Combined treatment of PEMF and osteo-induction medium increased ALP (A) and OC (B) mRNA expression. Experiments were performed three times with the similar results (n = 3 in each group). * indicates $P < 0.05$ vs Control, # indicates $P < 0.05$ vs PEMF and & indicates $P < 0.05$ vs OM. Abbreviations: Control, standard culture media; PEMF, PEMF exposure; OM, osteo-induction medium; PEMF + OM, combined treatments with PEMF and osteo-induction medium; ALP, alkaline phosphatase; OC, osteocalcin.




Figure 3 ALP activity was increased during the osteogenic differentiation of AECs. (A) The ALP activity was detected by BCIP/NBT staining (magnification of 400 \times). **(B)** The percent of ALP-positive cells (containing blue, insoluble, granular dye deposit) was calculated by ImageJ software. Experiments were performed three times with the similar results (n = 3 in each group). * indicates $P < 0.05$ vs Control, # indicates $P < 0.05$ vs PEMF and & indicates $P < 0.05$ vs OM.

expression of Nrf2 and Keap1. PEMF treatment alone significantly induced the expression of Nrf2 at day 7. Combined use of PEMF and osteo-induction medium evoked a significant up-regulation of Nrf2 mRNA expression compared with other treatments (Figure 2F). Furthermore, the gene expression profile of Keap1 was similar to that of Nrf2 during the osteogenic induction of AECs (Figure 6D,E).

Integrin β 1 may be the PEMF sensitive receptor in the process of osteogenic differentiation of AECs

Integrin β 1 is a well-known mechanoreceptor in mediating the transduction of mechanical strain in most cells [20]. We therefore investigated whether integrin β 1 might also act as the PEMF-sensitive receptor in the osteogenic induction of AECs. Real-time RT-PCR results showed that, PEMF alone induced the significant up-regulation of

gene expression of integrin β 1 in AECs at day 7. Moreover, treatments with PEMF and osteo-induction medium together evoked a significant up-regulation of integrin β 1 mRNA expression in comparison to the other treatments (Figure 6F). These results suggest that, integrin β 1 appears to be one of the PEMF or biochemical sensitive receptor mediated in the osteogenic differentiation of AECs.

Discussion

In the present study, the combined effects of physical treatment (PEMF) and biochemical stimuli (osteo-induction medium) on the osteogenic differentiation of AECs were investigated. The major findings of this study are: (1) Combined application of PEMF and osteo-induction medium to AECs has stronger effects on osteogenic differentiation, than either treatment alone; (2) Activation of signaling pathways, such as BMP-2 and Wnt/ β -catenin,




Figure 4 OC protein expression was induced to rise during the osteogenic differentiation of AECs. **(A)** The expression of OC protein was analyzed by immunocytochemistry (magnification of 400×). **(B)** The integrated optical density (IOD) of OC was measured by ImageJ software. Experiments were performed three times with the similar results (n = 3 in each group). * indicates $P < 0.05$ vs Control, # indicates $P < 0.05$ vs PEMF and & indicates $P < 0.05$ vs OM.

might be important in mediating the PEMF and/or osteo-induction medium-induced osteogenic differentiation of AECs; (3) Nrf2/Keap1, master regulators of ROS, might be implicated in the PEMF and/or osteo-induction medium-induced osteogenic differentiation of AECs; (4) The expression of integrin β 1 mRNA was up-regulated in the process of osteogenic differentiation of AECs induced by PEMF and/or osteo-induction medium. Our data indicate that PEMF could play an important role in the modulation of osteogenic differentiation of AECs. These observations also establish possible links of integrin β 1 and Nrf2/Keap1 in mediating PEMF and/or osteo-induction medium-induced osteogenic differentiation of AECs.

PEMF could interfere with cellular growth, proliferation and differentiation, as recently demonstrated in osteoblasts [9]. PEMF is also capable of regulating Ca^{2+} homeostasis and promoting fracture healing [21]. There have been several studies to demonstrate the inductive effect of PEMF in the osteoblast differentiation of MSCs [11,12,14,22,23]. Tsai et al. reported that PEMF could induce early onset of osteogenic differentiation of MSCs on the basis of ALP activity and stimulate the gene

expression of ALP and Runx-2 at day 7 but lower at day 10 in the process of osteogenic induction [22]. The similar results were reported by Sun et al. who found that exposure to PEMF significantly increased ALP gene expression during the early stages of osteogenesis and enhanced mineralization near the midpoint of osteogenesis [12]. Additionally, Song et al. revealed that PEMF could up-regulate the gene expression of Runx-2, bone sialoprotein (BSP) and osteopontin (OPN); enhance the alkaline phosphatase activity and calcium deposition in a time-dependent manner. Furthermore, MEK/ERK signaling pathway might be mediated in this process of osteogenic differences of MSCs [11].

The present study showed that PEMF stimulation alone could induce the expression of osteoblast markers ALP and OC at both gene and protein levels at specific time point(day 7). The osteo-induction medium was also able to induce the osteogenesis of AECs as reported in the previous literature [5]. Moreover, combined application of PEMF and osteo-induction medium led to significant up-regulation of ALP and OC expression and promoted the obvious extracellular matrix calcification.




Figure 5 A marked increase of calcium deposits was evoked during the osteogenic differentiation of AECs. (A) Calcium deposition was assessed by alizarin red S staining (arrows) (magnification of 400x). (B) The amount of calcified deposition was semi-quantitatively calculated with ImageJ software. Experiments were performed three times with the similar results (n = 3 in each group). * indicates $P < 0.05$ vs Control, # indicates $P < 0.05$ vs PEMF and & indicates $P < 0.05$ vs OM.

Our findings demonstrated the synergistic effects of physical (PEMF) and biochemical stimuli (OM) on the osteogenic induction of AECs. The effect of PEMF on the osteoblast differentiation of AECs may depend on the specific parameters of PEMF, such as waveform, duration, frequency and magnetic flux, as well as different cell types [22,24]. This may be the reason why the effects of PEMF alone on the stimulation of osteogenic differentiation of AECs were found most obvious at day 7. Therefore, further studies are necessary to determine the optical parameters of PEMF in the osteogenic differentiation of AECs.

Among the intracellular signals involved in PEMF actions, the activation of bone morphogenetic proteins (BMPs) plays important roles. Runx2, as a downstream regulator of BMP-2 signaling, is necessary for osteoblast differentiation [25]. Wnt/ β -catenin signaling is also of crucial importance for MSCs osteogenesis [26]. Our results showed that the gene expression of BMP-2, Runx2 and β -catenin were all up-regulated in the osteogenic differentiation of AECs induced by PEMF alone at the specific time point (day 7). Osteo-induction medium

alone or combined with PEMF exposure could induce BMP-2, Runx2 and β -catenin gene expression especially at the early stage of osteogenic induction (day 3). Additionally, similar gene expression profiles of BMP2, Runx2 and β -catenin were observed. These results may be due to the fact that both BMP-2/Runx2 and Wnt/ β -catenin signaling could play important role in activating the osteogenic induction of AECs at the early-stage, while down-regulation of these signals are required for the late-stage of osteogenesis and matrix mineralization [26,27]. Therefore, the present observations showed that PEMF and/or osteo-induction medium-induced the osteogenic differentiation of AECs may be via activation of both BMP-2 and Wnt/ β -catenin signaling.

Another pathway implicated in the PEMF action is the generation of reactive oxygen species (ROS) [28,29]. Recent study showed that Nrf2/Keap1, master regulator of ROS generation, would be required for intestinal stem cells (ISCs) proliferation [19]. In our study, the induction expression of Nrf2 and Keap1 were observed in the treatment of PEMF and/or osteo-induction medium, and the gene expression of Nrf2 and Keap1 exhibited similar




Figure 6 Molecular mechanisms linked to PEFM-induced osteogenic differentiation of AECs. Targeted genes involve: (A) bone morphogenetic protein 2 (BMP-2); (B) Runx-2; (C) β -catenin; (D) Nrf2; (E) Keap1; (F) Integrin β 1. Experiments were performed three times with the similar results (n = 3 in each group). * indicates $P < 0.05$ vs Control, # indicates $P < 0.05$ vs PEFM and & indicates $P < 0.05$ vs OM.

profiles during the osteogenesis of AECs. As key regulators of ROS generation, Nrf2/Keap1 might be of potential importance during osteogenic differentiation of AECs induced by PEFM and/or the osteo-induction medium.

The mechanism involving how the cells sense and transduce physical stimulation such as PEFM into biochemical signals has remained elusive. Integrins function as one of the mechanoreceptors, which are capable of switching mechanical strain to biochemical signals [20,30].

This process is comprised of binding to the extracellular matrix (ECM) ligands and activating the specific signaling pathways which would be involved in the mechanical-induced differentiation of cells [31]. Kasten et al. reported that certain biological functions of MSCs would be performed under the circumstance of integrin-mediated mechanical forces [32]. Franceschi et al. found that the application of mechanical force to osteoblasts could activate the MAPK signaling through integrin α 2 and β 1 [33]. However, little is known about the role of integrins in the

differentiation of AECs. The current study showed that, integrin β 1 gene expression was up-regulated in the PEMF and/or osteo-induction medium-induced osteogenic differentiation of AECs. We propose that integrin β 1 might act as a receptor, which can be inducible in response to the physical stimulation, especially to the PEMF.

Our preliminary study demonstrates the role of BMP-2, Wnt/ β -catenin, Nrf2/Keap1 and integrin β 1 in the osteogenic differentiation of AECs, only from the perspective of gene expression. Even though these molecules/pathways may be critical in mediating PEMF/osteo-induction medium enabled osteogenic differentiation, it is difficult to reveal the exact mechanisms without further testing, such as pathway-specific approaches. Therefore, additional experiments are needed to explore the mechanisms on how these signalings, ROS and integrin β 1 play role in regulating the PEMF-induced osteogenic differentiation of AECs.

Conclusions

The present study demonstrates that combined use of physical (PEMF) and biochemical stimuli (osteo-induction medium) is synergistic for the osteogenic differentiation of AECs, which might be a novel approach in bone regenerative medicine. BMP-2/Runx2, Wnt/ β -catenin, ROS and integrin signalings might be involved in the osteogenic differentiation of AECs induced by PEMF and/or osteo-induction medium.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

QW designed and carried out the experiments and performed statistical analysis and drafted the manuscript. Wc W designed the experiments and responsible for AEC cell isolation and culture; XY H worked on cell culture; A Z provided study material; J W and HQ C conception and design; S L worked on the transmission electron microscopy. CQ H conceived of the study. FM L participated in the design and coordination. XJ L conceived of the study, provided expert advice, interpretation of the study's results and prepared the manuscript. All authors read and approved the final manuscript.

Acknowledgments

This work was supported by grants from the National Natural Science Foundation of China, No. 30470437, 30870596 and 11072163 (XIAOJING LIU), No.81171865 (CHENGQI HE), No. 30971325(FENGMING LUO). The PEMF system was support by the Department of Rehabilitation Medicine, West China Hospital, Sichuan University. We would like to acknowledge the assistance and critical advice provided by Dr. Jue Lin (University of California, San Francisco) and Dr. Rui Lin (Medimmune Inc.) in the preparation of this manuscript.

Author details

¹Center of Rehabilitation Medicine, West China Hospital, Sichuan University, Chengdu, Sichuan, P. R. China. ²Rehabilitation Key Laboratory of Sichuan Province, Chengdu, Sichuan, P. R. China. ³Laboratory of Cardiovascular Diseases, Regenerative Medicine Research Center, West China Hospital, Sichuan University, Chengdu, Sichuan, P. R. China. ⁴Sichuan University-The Hong Kong Polytechnic University Institute for Disaster Management and Reconstruction, Sichuan University, Chengdu 610041, China. ⁵Department of Obstetrics and Gynecology, West China Second Hospital, Sichuan University, Chengdu, Sichuan, P. R. China. ⁶Department of Pathology, West

China Hospital, Sichuan University, Chengdu, Sichuan, P. R. China. ⁷Laboratory of Biomedical Engineering, West China Center of Medical Sciences, Sichuan University, Chengdu, Sichuan, P. R. China. ⁸Golden Card Ward, West China Hospital, Sichuan University, Chengdu, Sichuan, P. R. China.

Received: 17 February 2014 Accepted: 28 June 2014

Published: 11 August 2014

References

1. Miki T, Marongiu F, Ellis E: Isolation of amniotic epithelial stem cells. *Curr Protoc Stem Cell Biol* 2007, **12**:1-9.
2. Miki T, Mitamura K, Ross MA, Stolz DB, Strom SC: Identification of stem cell marker-positive cells by immunofluorescence in term human amnion. *J Reprod Immunol* 2007, **75**(2):91-96.
3. Carbone A, Paracchini V, Castellani S, Di Gioia S, Seia M, Colombo C, Conese M: Human Amnion-Derived Cells: Prospects for the Treatment of Lung Diseases. *Curr Stem Cell Res Ther* 2014, **9**(4):297-305.
4. Barboni B, Mangano C, Valbonetti L, Marruchella G, Berardinelli P, Martelli A, Muttini A, Mauro A, Bedini R, Turriani M: Synthetic bone substitute engineered with amniotic epithelial cells enhances bone regeneration after maxillary sinus augmentation. *PLoS One* 2013, **8**(5):e63256.
5. Parolini O, Alviano F, Bagnara GP, Bilic G, Buhring HJ, Evangelista M, Hennerbichler S, Liu B, Magatti M, Mao N: Concise review: isolation and characterization of cells from human term placenta: outcome of the first international Workshop on Placenta Derived Stem Cells. *Stem Cells* 2008, **26**(2):300-311.
6. H-f S, Xiong J, Chen Y-x, Wang J-f, Qiu X-s, Wang Y-h, Qiu Y: Early application of pulsed electromagnetic field in the treatment of postoperative delayed union of long-bone fractures: a prospective randomized controlled study. *BMC Musculoskelet Disord* 2013, **14**(1):35.
7. Assiotis A, Sachinis NP, Chalidis BE: Pulsed electromagnetic fields for the treatment of tibial delayed unions and nonunions. A prospective clinical study and review of the literature. *J Orthop Surg Res* 2012, **7**(1):1-6.
8. Vincenzi F, Targa M, Corciulo C, Gessi S, Merighi S, Setti S, Cadossi R, Goldring MB, Borea PA, Varani K: Pulsed electromagnetic fields increased the anti-inflammatory effect of A2A and A3 adenosine receptors in human T/C-28a2 chondrocytes and hFOB 1.19 osteoblasts. *PLoS One* 2013, **8**(5):e65561.
9. Barnaba S, Papalia R, Ruzzini L, Sgambato A, Maffulli N, Denaro V: Effect of pulsed electromagnetic fields on human osteoblast cultures. *Physiother Res Int* 2013, **18**(2):109-114.
10. Ceccarelli G, Bloise N, Mantelli M, Gastaldi G, Fassina L, Cusella De Angelis MG, Ferrari D, Imbriani M, Visai L: A Comparative analysis of the in vitro effects of pulsed electromagnetic field treatment on osteogenic differentiation of two different mesenchymal cell lineages. *Bio Res Open Access* 2013, **2**(4):283-294.
11. Song M-Y, Yu J-Z, Zhao D-M, Wei S, Liu Y, Hu Y-M, Zhao W-C, Yang Y, Wu H: The time-dependent manner of sinusoidal electromagnetic fields on rat bone marrow mesenchymal stem cells proliferation, differentiation, and mineralization. *Cell Biochem Biophys* 2014, **69**(1):47-54.
12. Sun LY, Hsieh DK, Lin PC, Chiu HT, Chiou TW: Pulsed electromagnetic fields accelerate proliferation and osteogenic gene expression in human bone marrow mesenchymal stem cells during osteogenic differentiation. *Bioelectromagnetics* 2010, **31**(3):209-219.
13. Bai W-f, Xu W-c, Feng Y, Huang H, Li X-p, Deng C-y, Zhang M-s: Fifty-Hertz electromagnetic fields facilitate the induction of rat bone mesenchymal stromal cells to differentiate into functional neurons. *Cytotherapy* 2013, **15**(8):961-970.
14. Zhong C, Zhang X, Xu Z, He R: Effects of low-intensity electromagnetic fields on the proliferation and differentiation of cultured mouse bone marrow stromal cells. *Phys Ther* 2012, **92**(9):1208-1219.
15. Yang F, Yang D, Tu J, Zheng Q, Cai L, Wang L: Strontium enhances osteogenic differentiation of mesenchymal stem cells and in vivo bone formation by activating wnt/catenin signaling. *Stem Cells* 2011, **29**(6):981-991.
16. Pan P, Fu H, Zhang L, Huang H, Luo F, Wu W, Guo Y, Liu X: Angiotensin II upregulates the expression of placental growth factor in human vascular endothelial cells and smooth muscle cells. *BMC Cell Biol* 2010, **11**(1):36-44.
17. Dalby MJ, Gadegaard N, Tare R, Andar A, Riehle MO, Herzyk P, Wilkinson CDW, Oreffo ROC: The control of human mesenchymal cell differentiation using nanoscale symmetry and disorder. *Nat Mater* 2007, **6**(12):997-1003.

18. Yanes O, Clark J, Wong DM, Patti GJ, Sanchez-Ruiz A, Benton HP, Trauger SA, Despons C, Ding S, Siuzdak G: **Metabolic oxidation regulates embryonic stem cell differentiation.** *Nat Chem Biol* 2010, **6**(6):411–417.
19. Hochmuth CE, Biteau B, Bohmann D, Jasper H: **Redox regulation by Keap1 and Nrf2 controls intestinal stem cell proliferation in *Drosophila*.** *Cell Stem Cell* 2011, **8**(2):188–199.
20. Puklin-Faucher E, Sheetz MP: **The mechanical integrin cycle.** *J Cell Sci* 2009, **122**(2):179–186.
21. Ibiwoye MO, Powell KA, Grabiner MD, Patterson TE, Sakai Y, Zborowski M, Wolfman A, Midura RJ: **Bone mass is preserved in a critical-sized osteotomy by low energy pulsed electromagnetic fields as quantitated by in vivo micro-computed tomography.** *J Orthop Res* 2004, **22**(5):1086–1093.
22. Tsai MT, Li WJ, Tuan RS, Chang WH: **Modulation of osteogenesis in human mesenchymal stem cells by specific pulsed electromagnetic field stimulation.** *J Orthop Res* 2009, **27**(9):1169–1174.
23. Yang Y, Tao C, Zhao D, Li F, Zhao W, Wu H: **EMF acts on rat bone marrow mesenchymal stem cells to promote differentiation to osteoblasts and to inhibit differentiation to adipocytes.** *Bioelectromagnetics* 2010, **31**(4):277–285.
24. Schwartz Z, Simon B, Duran M, Barabino G, Chaudhri R, Boyan B: **Pulsed electromagnetic fields enhance BMP2 dependent osteoblastic differentiation of human mesenchymal stem cells.** *J Orthop Res* 2008, **26**(9):1250–1255.
25. Satija NK, Gurudutta G, Sharma S, Afrin F, Gupta P, Verma YK, Singh VK, Tripathi R: **Mesenchymal stem cells: molecular targets for tissue engineering.** *Stem Cells Dev* 2007, **16**(1):7–24.
26. Ling L, Nurcombe V, Cool SM: **Wnt signaling controls the fate of mesenchymal stem cells.** *Gene* 2009, **433**(1–2):1–7.
27. Horst G, Werf SM, Farih-Sips H, Bezooijen RL: **Downregulation of Wnt signaling by increased expression of dickkopf-1 and -2 is a prerequisite for late-stage osteoblast differentiation of KS483 cells.** *J Bone Miner Res* 2005, **20**(10):1867–1877.
28. Regoli F, Gorbi S, Machella N, Tedesco S, Benedetti M, Bocchetti R, Notti A, Fattorini D, Piva F, Principato G: **Pro-oxidant effects of extremely low frequency electromagnetic fields in the land snail *Helix aspersa*.** *Free Radic Biol Med* 2005, **39**(12):1620–1628.
29. Frahm J, Mattsson MO, Simko M: **Exposure to ELF magnetic fields modulate redox related protein expression in mouse macrophages.** *Toxicol Lett* 2010, **192**(3):330–336.
30. Ward DF Jr, Salasnyk RM, Klees RF, Backiel J, Agius P, Bennett K, Boskey A, Plopper GE: **Mechanical strain enhances extracellular matrix-induced gene focusing and promotes osteogenic differentiation of human mesenchymal stem cells through an extracellular-related kinase-dependent pathway.** *Stem Cells Dev* 2007, **16**(3):467–480.
31. Yim E, Sheetz MP: **Force-dependent cell signaling in stem cell differentiation.** *Stem Cell Res Ther* 2012, **3**:41.
32. Kasten A, Müller P, Bulnheim U, Gröll J, Bruellhoff K, Beck U, Steinhoff G, Möller M, Rychly J: **Mechanical integrin stress and magnetic forces induce biological responses in mesenchymal stem cells which depend on environmental factors.** *J Cell Biochem* 2010, **111**(6):1586–1597.
33. Franceschi RT, Xiao G: **Regulation of the osteoblast-specific transcription factor, Runx2: Responsiveness to multiple signal transduction pathways.** *J Cell Biochem* 2003, **88**(3):446–454.

doi:10.1186/1471-2474-15-271

Cite this article as: Wang et al.: Osteogenic differentiation of amniotic epithelial cells: synergism of pulsed electromagnetic field and biochemical stimuli. *BMC Musculoskeletal Disorders* 2014 **15**:271.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

