



Received: 28 February 2019  
Accepted: 20 May 2019  
First Published: 29 May 2019

\*Corresponding author: Xuemei Bai,  
Institute of Life Science and  
Technology, ENN Group, Langfang,  
065001, China  
E-mail: [xuemei\\_bai@yahoo.com](mailto:xuemei_bai@yahoo.com)

Reviewing editor:  
Tianzhu Zang, Systems  
Pharmacology & Translational  
Therapeutics, University of  
Pennsylvania Perelman School of  
Medicine, Philadelphia, PA, USA

Additional information is available at  
the end of the article

## CELL, MOLECULAR & DEVELOPMENTAL BIOLOGY | RESEARCH ARTICLE

# Evaluation of the extremely low-frequency electromagnetic field (ELF-EMF) on the growth of bacteria *Escherichia coli*

Yu Chen<sup>1</sup>, Zhongzhen Cai<sup>1</sup>, Qian Feng<sup>1</sup>, Peng Gao<sup>1</sup>, Yongdong Yang<sup>1</sup>, Xuemei Bai<sup>1\*</sup> and Bruce Q. Tang<sup>1</sup>

**Abstract:** The goal of this work was to evaluate the biological effect of the extremely low frequencies (ELFs). We started by evaluating the effect of frequencies with the anti-bacterial functions listed in the Rife frequency database on the growth of *Escherichia coli*. Among the 11 sets of frequencies tested, we found that only 1 set comprising frequencies below 100 Hz showed a significant effect on the growth of *E. coli*, and it was an enhancement effect instead of an inhibition on the growth. We further tested various sets of frequencies without anti-bacterial functions but comprising frequencies below 100 Hz, and they all exhibited a significant promoting effect on the growth of *E. coli* when exposed to the irradiation for 1 h. We further evaluated the effect of the ELF on the growth of *E. coli* by scanning from 0 to 100 Hz. The results showed that the growth rate was noticeably enhanced for the group exposed to the irradiation at the range of 0–100 Hz, with the range of 40–60 Hz being the most effective. In addition, the effect was time dependent, demonstrated by a significant increase of the growth rate observed after exposure of 1 h but a slight decrease after exposure of 16 h. The results of this study provided robust data on the biological effect of ELF on the growth of *E. coli*, an important group of

### ABOUT THE AUTHOR

Xuemei Bai received a PhD and postdoctoral training in Biological Sciences from USM and USC, specializing in biotechnologies including biofuels and novel bio-products production. Her current research interests include energy medicine and biological medicine, mainly looking into the effects of physical fields and qi-invigorating medicine on the health and physiological processes of biological systems.

### PUBLIC INTEREST STATEMENT

This study reported the significant biological effect of extremely low-frequency electromagnetic irradiation on the growth of *E. coli*. The results demonstrated that the growth rate was noticeably affected when treated by the EMI of frequencies below 100 Hz, whether they are a combination of frequencies below 100 Hz or scanning frequencies in the range of 0–100 Hz. It was found that the frequency in the range of 40–60 Hz had the most significant effect on the growth rate. The EMI effect of 50 Hz on the growth of bacteria was time dependent, with an increase in growth within 2 h of irradiation, and an inhibition when the exposure time was over 4 h. These findings have applicable values for the fine chemical fermentation processes as the biomass of *E. coli* could be enhanced significantly through the appropriate EMI stimulation. On the other hand, it also raises concerns regarding the safety of these frequencies on the biological systems.

human microbiota, which has critical implication on the safety risk of these frequencies on human health.

**Subjects: Medicinal & Pharmaceutical Chemistry; Environmental Chemistry; Medical Physics; Biomedical Engineering; Electromagnetics & Communication; Electronic Devices & Materials**

**Keywords: extremely low-frequency electromagnetic field (ELF-EMF); electromagnetic irradiation (EMI) effects; bacterial growth; Escherichia coli**

## 1. Introduction

In the natural world, almost all living organisms are “immersed” in a variety of electromagnetic field (EMF) and interact with them. Living organisms including bacteria have great diversity and may have different sensitivity to EMF, which are known to have wide spectrum and different characteristics (Soghomonyan, Trchounian, & Trchounian, 2016). For many years, the electromagnetic irradiation (EMI) effects of the extremely low-frequency electromagnetic field (ELF-EMF) on biological systems have been debated. Some studies showed there was no significant biological effect (De Haas, Lazarovici, & Morrison, 1979; Devereaux, Hazleman, & Thomas, 1985; Harrison & Bassett, 1997; Karasek, Czernicki, & Woldanska-Okonska et al., 2000), while others demonstrated significant effect of the ELF irradiation (Amaroli et al. 2006; Akbarnejad et al., 2017; Gu, Lu, & Wu et al., 2012; Mittenzwey, Sussmuth, & Mei, 1996; Rodriguez Justo, Perez, & Chacon Alvarez et al., 2006), especially the frequency of 50 Hz, which can be found everywhere in daily life (such as power lines, ordinary household appliances, and communication devices) (Simko & Mattsson, 2004). As for the biological effect on human health, both harmful and useful effects have been reported (Feychting, Forssén, & Floderus, 1997; Li, Thériault, & Lin, 1997; Saffer & Thurston, 1995; Savitz, Wachtel, Barnes, John, & Tvrdik, 1988; Wertheimer & Leeper, 1979). The root cause of the conflicting results is due to the lack of essential controls for the experiments conducted (Saffer & Thurston, 1995).

The ELF-EMF effects are dependent on various factors, such as frequency, field strength, exposure time, cultivation mode, and genotype (Fojt, Strasak, & Vetterl et al., 2004; Strasak, Vetterl, & Smarda, 1998, 2002). Most work with cellular systems was performed by choosing one specific frequency or one modulation frequency with varying field intensities. Screening of a wide range of frequency and intensity has not yet been done.

As for the mechanism of the effect, water, as the major constituent of growth medium as well as biological systems, was speculated to play a role in mediating the EMI effect on *E. coli*, and the mediated effect through water at extremely high frequencies of 1.8, 53, 70.6, and 73 GHz on *E. coli* growth was clearly demonstrated (Torgomyan, Hovnanyan, & Trchounian, 2013; Torgomyan, Kalantaryan, & Trchounian, 2011). However, there is no information regarding the mediating effect of water for ELF.

For this study, *Escherichia coli* strain DH5 $\alpha$  was chosen to examine the EMI effect of ELF. We started by examining numerous frequency combinations with anti-bacterial functions (i.e. *E. coli* inhibitory programs, chlorine, ozone, antibiotic and general antiseptic programs) listed in the Rife database (<https://www.spooky2.com/>), which is a compilation of frequencies from Rife frequency researchers and can be accessed and downloaded online, and then focused on the frequency combinations below 100 Hz.

To better understand the effect of the ELF, we designed an ELF exposure system to generate a weak uniform ELF-EMF. By using this system, a screening study was performed by sweeping frequency in the range of 0–10,000Hz initially, then narrowed down to the range of 0–100 Hz, and followed by focusing on the frequency of 50 Hz. The primary goal was to gain more knowledge of frequency that showed the most significant impact on the growth of *E. coli*.

By simultaneously examining the effect of different ranges of frequencies, including scanning a few distinctive frequencies or sweeping in different ranges of frequencies, it increased the robustness of the results compared with looking into one frequency only. In addition, with the effort to shed light on the mechanism of biological effect, we also examined the mediating effect of 50 Hz through water and growth media on the growth of *E. coli*.

## 2. Materials and methods

### 2.1. Bacteria and preparation

*Escherichia coli* strain DH5 (obtained from the Institute of ENNOVA Life Science and Technology) was maintained on Luria-Bertani (LB) medium with 1.5% agar (Cat# A8190, Solarbio) at 4°C in the freezer.


For each experiment, it was taken from freezer and suspended in a 50 mL Erlenmeyer flask containing 30 mL of LB liquid medium, comprising 5 g/L yeast extract (Cat#: LP0021, Oxoid), 10 g/L Tryptone (Cat#: LP0042, Oxoid), 10 g/L NaCl (Cat#: S5886-1KG, Sigma), with pH set to 7, and agitation of 200 rpm on a rotary shaker at 37°C for 12 h. This process ensured that bacteria were activated into the stage of exponential growth, thereafter the activated cells were used as a seed of inoculation for subsequent irradiation experiments.

For the irradiation experiments, the seed was inoculated into LB medium to obtain the suspensions of bacterial cells with an optical density of OD<sub>600 nm</sub> = 0.1 (equivalent to a cell density of  $1.0 \times 10^6 \sim 10^7$  cells per mL suspension). Then, the suspension was transferred into 50 mL Erlenmeyer flasks containing 5 mL of suspension (thickness of the suspension was about 5 mm) to be used as the test samples. It should be noted that the cell density and thickness were optimized in the foregoing experiments to make sure the test sample was prepared appropriately for all experiments.

### 2.2. EMI procedure

Figure 1 shows the schematic diagram of the ELF exposure system. ELF exposure system including coils and culture flask has been specifically designed so it can fit inside the incubator with chambers to keep the culture flasks in place. The samples were divided into two groups, with one as the EMF-treated group and the other the control or untreated group. In order to make sure the ELF-EMF generated was uniform for all samples, a flat coil (47 mm diameter, 7 mm height) was adopted to fit the bottom of the culture flask. The irradiation procedure was performed by connecting the coil to a commercially available frequency generator (FY2300 type, FeelTech Enterprise, China, <http://www.feeltech.net/>).

**Figure 1. Schematic illustration of the experimental design used for the measurement of growth characteristics of *E. coli*.**



- (i) For evaluation, the EMI effect of Rife frequencies (<https://www.spooky2mall.com/download/spooky2rifefrequencylist.pdf>) on *E. coli*, 3 groups and a total of 14 programs of frequencies were used for the experiment (Table 1). Group A included 7 programs which were related to the inhibitory effect of *E. coli*, group B encompassed 4 programs regarding frequencies with anti-bacterial properties, and group C included programs with frequencies below 100 Hz.
- (ii) For the wave-scanning effect experiment, the scanning wave was set in the ranges of 0–100, 100–500, 500–1000, 0–1000, and 0–10,000 Hz, respectively; for frequency lower than 100 Hz, the scanning wave was set in the ranges of 0–20, 20–40, 40–60, 60–80, and 80–100 Hz, respectively. The duration of irradiation was set for 1 h.
- (iii) For the single-wave effect experiment, a frequency of 50 Hz was chosen to examine the EMI effect with respect to the exposure time. The irradiation time was set for 1, 2, 4, 8, and 16 h, and all the other variables were kept the same as the wave-scanning effect experiment.
- (iv) For investigating the mediated effect of the frequency at 50 Hz on bacteria, three sets of samples were set up, which included (1) growth medium with bacterial culture (MB), (2) growth medium only (M), and (3) double-distilled water only (W). For the EMF-treated groups, the three samples were exposed to the 50 Hz irradiation at room temperature for 1 h without agitation. After irradiation, fresh medium was added into MB, bacteria (same batch as used for MB) into M, and bacteria (same batch as used for MB) and the growth medium into W, followed by adjusting all the samples to the appropriate cell density as stated above. Upon finishing, all samples were immediately placed into the incubator at 37°C with agitation of 120 rpm for 1 h.

**Table 1. List of Rife frequency used for evaluation of EMI effect on *E. coli***

Groups	Programs*	Frequencies (Hz)	Description
A	<i>E. coli</i> 1 (CAFL)	332.5, 798, 1729, 7847	Clearly related to the inhibitory effect of <i>Escherichia coli</i> .
	<i>E. coli</i> 1 (XTRA)	971.66, 974.14, 12,250, 12,281.25,	
	<i>E. coli</i> 3 (XTRA)	882.44, 11,125,	
	<i>E. coli</i> 5 (XTRA)	330, 358, 539, 556, 634, 642, 776, 800, 840, 856, 934, 1000, 1244, 1712, 1730, 7847,	
	<i>E. coli</i> 6 (XTRA)	882.44, 974.14, 17,724.2, 19,566.31,	
	<i>E. coli</i> (HC)	356,000,	
	<i>E. coli</i> (BIO)	282, 333, 413, 957, 1320, 1722,	
B	Wide Spectrum Antibiotic 4 (XTRA)	727, 787, 802, 880, 465,	Related to the general inhibition of bacteria.
	Chlorine 35cl (XTRA)	208.56, 224.84, 19,136.72,	
	Chlorine 37cl (XTRA)	173.62, 187.15, 15,929.69,	
C	Ozone Generate (XTRA)	78, 16, 15.99,	Not related to the biological effects of bacteria, but with similar to frequencies of "Ozone Generate"
	Accelerate Injury Healing (XTRA)	47,	
	Adynamia Geriatric (XTRA)	60, 27.5,	
	Frequency Fatigue (CAFL)	10.55, 7.83,	

\*CAFL, XTRA, HC, and BIO are the sub-databases from the Rife frequency list. CAFL is the Consolidated Annotated Frequency List, amassed from the experience of Rife experimenters over years. XTRA is a collection of sets from various sources, all chosen for their reputation for effectiveness. HC is Dr. Hulda Clark's database. BIO is based on Russian frequency research.

For the Rife frequencies evaluation experiment, a 10 V wave scanning with different programs of Rife frequencies (intensity range: 0.1–3.0 mT) was used. For the wave-scanning effect experiment, a 10 V sine wave scanning was generated in the frequency range of 0–10,000 Hz (0.1–3.0 mT) with the repetition rate of 60 s, and for the single-wave effect experiment, a continuous 20 V sine wave at the frequency of 50 Hz (2.3–2.5 mT) was generated.

To investigate the growth characteristics of *E. coli* exposed to the ELF-EMF, the temperature was set at 37°C in an incubator with the temperature variation less than 0.3°C (non-thermal effects). The samples were kept in suspension by placing on a shaker at 120 rpm during the experiment.

For all experiments, the control, also referred to as the untreated sample, was set up in triplicates under the same culturing conditions (e.g. placed in the same type of coil in the same incubator) as that for the treated group. The only difference is that the control group received no irradiation by keeping the frequency generator powered off.

### 2.3. Bacterial growth determination

The growth of cells was indicated by monitoring optical density at a wavelength of 750 nm using a UV-VIS Auto spectrophotometer (UV-2800 type, UNIC, China). The growth rate of culture was calculated as the optical density change over time ( $R = \frac{OD_T - OD_0}{T}$ ) by taking OD of samples at the time of inoculation (0) and at the end of the exposure (T). EMF exposure effect was assessed in terms of the variation of growth rate between the treated and the untreated group. The degree of variation was calculated as follows:  $V(\%) = \frac{R_t - R_u}{R_u}$ , where  $R_t$  and  $R_u$  were the growth rates of the treated and the untreated groups. Since the only difference between the treated and untreated groups was the irradiation, the variation of growth rate between these two groups is used to indicate the effect of the irradiation. All experiments were performed in triplicates with the data averaged and their standard deviation calculated.

## 3. Results

### 3.1. EMI effect of Rife frequencies on bacterial growth

Three groups of programs were tested. The results showed that the growth rate was not affected substantially under the irradiation of the frequency programs from group A (Figure 2(a)). Three sets of frequencies tested enhanced the growth slightly, while the other four sets suppressed the growth, with effect less than 7%. For the effect of programs of group B, there was a significant enhancement induced by the frequencies of “Ozone Generate”, and the increase in growth reached 25%, while the effect with other programs was less than 10%. The Ozone Generate program contains three frequencies under 100 Hz (78, 16, 15.99 Hz), so we further examined programs in group C with frequencies below 100 Hz but not related to anti-bacterial functions. The results showed a significant enhancement in growth by all programs of group C (Figure 2(b)).


### 3.2. EMI effect on bacterial growth for frequencies at different ranges

The growth rate was not affected when scanning in a relatively large range of frequency such as 0–10,000 and 0–1,000 Hz. However, the growth rate was slightly enhanced at the lower range of frequencies when scanning at a relatively smaller range such as 500–1,000 and 100–500 Hz; by lowering the frequency to 0–100 Hz, the growth rate was significantly enhanced with an average greater than 10% (Figure 3(a)). By narrowing down the range of the frequency to 0–100 Hz, the EMI effect on *E. coli* growth was much more pronounced with the maximum increase reaching over 20% (Figure 3(b)). The effect was frequency dependent, and the ranges of 40–60 and 60–80 Hz were the most effective in promoting the growth of *E. coli*. Based on these results, the frequency of 50 Hz was selected for subsequent single-wave experiments.

### 3.3. EMI effects on bacterial growth with respect to exposure time

The EMI effect for frequency at 50 Hz was dependent on exposure time as shown in Figure 4, where a significant increase in growth rate (>25%) was observed for groups with exposure time of

**Figure 2. The growth rate variation of *E. coli* after irradiation by EMF at different Rife frequencies programs: (a) programs from group A related to the inhibitory effect of *E. coli* and group B with anti-bacterial properties (b) Programs from group C with frequencies below 100 Hz, with no anti-bacterial properties stated in the database. Error bars representing standard deviation of triplicate samples.**



1 and 2 h, while no promotion in growth was observed for groups with exposure time of 4, 8, and 16 h. On the other hand, the prolonged exposure time (8 and 16 h) resulted in a slight decrease in growth rates (6%) for the treated group relative to the untreated group.


### 3.4. EMI effect on bacteria and indirect effect through medium and water

Consistent with the results obtained above, as shown in Figure 5, there was a significant increase in growth rate for the group with both the growth medium and bacteria being exposed to the 50 Hz irradiation for 1 h. However, for the other groups, no difference in growth rates was observed between the treated and untreated groups when the growth medium or the water was exposed to the 50 Hz irradiation only. Therefore, the EMI effect was only observed when bacteria were directly exposed to the 50 Hz frequency, and no effect in growth was obtained by simply treating the media or water with the 50 Hz frequency in this experiment.


## 4. Discussion

Some of the frequency combinations supposed to produce an inhibition effect in the Rife database exhibited a promoting effect on the growth of *E. coli*, especially the “Ozone Generate” program containing three frequencies below 100 Hz. So we further selected other programs with

**Figure 3.** The growth rate variation of *E. coli* after irradiation by EMF at different frequency ranges: (a) the scanning wave in the range of 0–10,000 Hz, and (b) the scanning wave in the range of 0–100 Hz. Control: EMF-treated with 0 Hz radiation. Error bars representing standard deviation of triplicate samples.




**Figure 4.** The EMF effect of frequency at 50 Hz on the growth rate of *E. coli* at different exposure time: 1, 2, 4, 8, and 16 h. Control: EMF-treated with 0 Hz radiation for 1 and 16h, respectively. Error bars representing standard deviation of triplicate samples.



frequencies below 100 Hz, but with no anti-bacterial functions stated in the database, and found the same enhancing effect on the growth of *E. coli*. This is a very intriguing result which led us to systematically examine the effect of frequencies under 100 Hz on the growth of *E. coli*.

**Figure 5.** The EMF effect of frequency at 50 Hz on *E. coli* growth rate through irradiation of medium with bacteria (MB), medium only (M), and water only (W). Error bars representing standard deviation of triplicate samples.



The issue of health hazards associated with EMFs produced by power lines or household appliances has been continuously debated, including different effects and the mechanism of these effects. Some even argued that such an effect may or may not exist due to the lack of essential controls of many researches published in this area (Saffer & Thurston, 1995). To avoid such a concern, we used frequency at 0 Hz as control and examined multiple ranges of frequencies simultaneously, which can all be compared to the targeting frequency. The results obtained here provided direct evidence of biological effects exerted by ELF-EMI at 0–100 Hz, especially 50 Hz.

It was reported that the cell cycle of human peripheral lymphocytes in culture was accelerated when exposed to 50 Hz irradiation of 5 mT (Antonopoulos, Yang, Stamm, Heller, & Obe, 1995), while an increased cell proliferation is regarded as cell transformation (Katsir & Parola, 1998). A growth increase of anchorage-independent JB6 cells after exposure to a 60 Hz (1.1 mT) was demonstrated (West, Hinson, Lyle, & Swicord, 1994). The EMI effect of 50 Hz on bacteria is of special interest because EMFs (50/60 Hz) can be caused by an alternating current which is inevitable for modern society.

In addition, EMI effect of 50 Hz showed the exposure time dependence, where a significant increase in the growth after exposure for 1 h but a slight decrease after exposure for 4 h were observed. More in-depth study is needed to address the reasons for the time-dependent effect. It could be cell damage due to EMI exposure over time or it could be caused by worse and worse growth conditions in the treatment group due to the fast growth in the first couple of hours. Therefore, to evaluate the EMI effect, the cell density, cell viability, and parameters of the growth media should be all taken into consideration.

Many investigations suggested that EMI is absorbed by water and the effects on bacteria can be mediated by the changes in water cluster structures (Binhi & Rubin, 2007; Fesenko, Geletyuk, Kazachenko, & Chemeris, 1995; Suresh, Satish, & Choudhary, 2006), and these effects might be long term (Belovolova, Glushkov, Vinogradov, Babintsev, & Golovanov, 2009; Fesenko et al., 1995; Sunkari, Aranovitch, Portwood, & Nikoshkov, 2011). However, unlike the EHF-EMF, water or media treated with ELF-EMF did not induce an EMI effect for bacteria cultured with it, indicating no mediated effect (indirect effect) of ELF-EMF for this study. For future work, the experiment should be performed in higher EMF levels to investigate whether stronger ELF-EMF is effective for a mediating effect through medium or water.



## 5. Conclusions

This study reported the significant biological effect of ELF-EMI on the growth of *E. coli*. The results demonstrated that the growth rate was noticeably affected when treated by the EMI of frequencies below 100 Hz, whether they are a combination of frequencies below 100 Hz or scanning frequencies in the range of 0–100 Hz. It was found that the frequency in the range of 40–60 Hz had the most significant effect on the growth rate. The EMI effect of 50 Hz on the growth of bacteria was time dependent, with an increase in growth within 2 h of irradiation, and an inhibition when the exposure time was over 4 h. It was demonstrated that the 50 Hz irradiation was a direct effect on the bacteria cells as treating media or water alone did not yield an EMI effect for bacteria cultured with it. These findings have applicable values for the fine chemical fermentation processes as the biomass of *E. coli* could be enhanced significantly through the appropriate EMI stimulation. On the other hand, it also raises concerns regarding the safety of these frequencies on the biological systems, which needs to be examined more systematically.

### Author Statement

The authors at ENN Group are working to establish an energy medicine system bridging the traditional Chinese medicine and modern western medicine. We are interested in the relationship between mitochondrial function and qi, as well as the effect of macroscopic entanglement of biological information. We are also interested in the biological effects induced by the physical fields (e.g. the electromagnetic, scalar, and torsion), as well as the non-local biological effects of molecules transferred by such physical fields.

### Acknowledgements

The authors would like to thank the ENN Group for the funding.

### Funding

This work was supported by the ENN Group [201908];

### Author details

Yu Chen<sup>1</sup>  
E-mail: [chenyuq@enn.cn](mailto:chenyuq@enn.cn)  
Zhongzhen Cai<sup>1</sup>  
E-mail: [caizhongzhen@enn.cn](mailto:caizhongzhen@enn.cn)  
Qian Feng<sup>1</sup>  
E-mail: [fengqian@enn.cn](mailto:fengqian@enn.cn)  
Peng Gao<sup>1</sup>  
E-mail: [gaopeng@enn.cn](mailto:gaopeng@enn.cn)  
Yongdong Yang<sup>1</sup>  
E-mail: [yangyongdong@enn.cn](mailto:yangyongdong@enn.cn)  
Xuemei Bai<sup>1</sup>  
E-mail: [xuemei\\_bai@yahoo.com](mailto:xuemei_bai@yahoo.com)  
Bruce Q. Tang<sup>1</sup>  
E-mail: [tangqing@enn.cn](mailto:tangqing@enn.cn)  
<sup>1</sup> Institute of Life Science and Technology, ENN Group, Langfang 065001, China.

### Competing Interests

The authors declare no conflict of interest.

### Citation information

Cite this article as: Evaluation of the extremely low-frequency electromagnetic field (ELF-EMF) on the growth of bacteria *Escherichia coli*, Yu Chen, Zhongzhen Cai, Qian Feng, Peng Gao, Yongdong Yang, Xuemei Bai & Bruce Q. Tang, *Cogent Biology* (2019), 5: 1625104.

### References

Akbarnejad, Z., Eskandary, H., Vergallo, C., Nematollahi-Mahani, S. N., Dini, L., Darvishzadeh-Mahani, F., & Ahmadi, M. (2017). Effects of extremely low-frequency pulsed electromagnetic fields (ELF-PEMFs) on glioblastoma cells (U87).

- Electromagnetic Biology and Medicine*, 36, 238–247. doi:10.1080/15368378.2016.1251452
- Amaroli, A., Trielli, F., Bianco, B., Giordano, S., Moggia, E., & Corrado, M. U. D. (2006). Effects of a 50 Hz magnetic field on dictyostelium discoideum (Protista). *Bioelectromagnetics*, 27, 528–534. doi:10.1002/bem.20240
- Antonopoulos, A., Yang, B. C., Stamm, A., Heller, W.-D., & Obe, G. (1995). Cytological effects of 50-hz electromagnetic-fields on human-lymphocytes in-vitro. *Mutation Research Letters*, 346, 151–157. doi:10.1016/0165-7992(95)90047-0
- Belovolova, L. V., Glushkov, M. V., Vinogradov, E. A., Babintsev, V. A., & Golovanov, V. I. (2009). Ultraviolet fluorescence of water and highly diluted aqueous media. *Physics of Wave Phenomena*, 17, 21–31. doi:10.3103/S1541308X0901004X
- Binhi, V. N., & Rubin, A. B. (2007). Magnetobiology: The kT paradox and possible solutions. *Electromagnetic Biology and Medicine*, 26, 45–62. doi:10.1080/15368370701205677
- De Haas, W. G., Lazarovici, M. A., & Morrison, D. M. (1979). The effect of low frequency magnetic fields on the healing of the osteotomized rabbit radius. *Clinical Orthopaedics and Related Research*, 145, 245–251.
- Devereaux, M. D., Hazleman, B. L., & Thomas, P. P. (1985). Chronic lateral humeral epicondylitis—A double-blind controlled assessment of pulsed electromagnetic field therapy. *Clinical and Experimental Rheumatology*, 3, 333–336.
- Fesenko, E. E., Geletyuk, V. I., Kazachenko, V. N., & Chemeris, N. K. (1995). Preliminary microwave irradiation of water solutions changes their channel-modifying activity. *FEBS Letters*, 366, 49–52.
- Feychting, M., Forssén, U., & Floderus, B. (1997). Occupational and residential magnetic field exposure and leukemia and central nervous system tumors. *Epidemiology*, 8, 384–389.
- Fojt, L., Strasak, L., Vetterl, V., & Šmarda, J. (2004). Comparison of the low-frequency magnetic field effects on bacteria *Escherichia coli*, *Leclercia ade-carboxylata* and *Staphylococcus aureus*. *Bioelectrochemistry*, 63, 337–341. doi:10.1016/j.bioelechem.2003.11.010
- Gu, S., Lu, G., Wu, Y., Li, S., Zhao, Y., & Li, K. (2012). *A study of the interaction between ELF-EMF and bacteria*. Springer, Berlin, Heidelberg.
- Harrison, M. H. M., & Bassett, C. A. L. (1997). The results of a double-blind trial of pulsed electromagnetic frequency in the treatment of Perthes' disease. *Journal of Pediatric Orthopaedics*, 17, 264–265.
- Karasek, M., Czernicki, J., Woldanska-Okonska, M., Zylinska, K., Swietoslowski, J. (2000). Chronic

- exposure to 25–80- $\mu$ T, 200-Hz magnetic field does not influence serum melatonin concentrations in patients with low back pain. *Journal of Pineal Research*, 29, 81–85.
- Katsir, G., & Parola, A. H. (1998). Enhanced proliferation caused by a low frequency weak magnetic field in chick embryo fibroblasts is suppressed by radical scavengers. *Biochemical and Biophysical Research Communications*, 252, 753–756. doi:10.1006/bbrc.1998.9579
- Li, C.-Y., Thériault, G., & Lin, R. S. (1997). Residential exposure to 60-Hertz magnetic fields and adult cancers in Taiwan. *Epidemiology*, 8, 25–30. doi:10.1097/00001648-199701000-00004
- Mittenzwey, R., Sussmuth, R., & Mei, W. (1996). Effects of extremely low-frequency electromagnetic fields on bacteria—The question of a co-stressing factor. *Bioelectrochemistry and Bioenergetics*, 40, 21–27. doi:10.1016/0302-4598(95)00504-8
- Rodriguez Justo, O., Perez, V. H., Chacon Alvarez, D., Monte Alegre, R. (2006). Growth of *Escherichia coli* under extremely low-frequency electromagnetic fields. *Applied Biochemistry and Biotechnology*, 134, 155–163.
- Saffer, J. D., & Thurston, S. J. (1995). Cancer risk and electromagnetic fields. *Nature*, 375, 22. doi:10.1038/375196b0
- Savitz, D. A., Wachtel, H., Barnes, F. A., John, E. M., & Tvrđik, J. G. (1988). Case-control study of childhood cancer and exposure to 60-hz magnetic fields. *American Journal of Epidemiology*, 128, 21–38. doi:10.1093/oxfordjournals.aje.a114943
- Simko, M., & Mattsson, M. O. (2004). Extremely low frequency electromagnetic fields as effectors of cellular responses in vitro: Possible immune cell activation. *Journal of Cellular Biochemistry*, 93, 83–92. doi:10.1002/jcb.20198
- Soghomonyan, D., Trchounian, K., & Trchounian, A. (2016). Millimeter waves or extremely high frequency electromagnetic fields in the environment: What are their effects on bacteria? *Applied Microbiology and Biotechnology*, 100, 4761–4771. doi:10.1007/s00253-016-7538-0
- Strasak, L., Vetterl, V., & Smarda, J. (1998). The effect of low-frequency electromagnetic fields on living organisms. *Sbornik Lekarsky*, 99, 455–464.
- Strasak, L., Vetterl, V., & Smarda, J. (2002). Effects of low-frequency magnetic fields on bacteria *Escherichia coli*. *Bioelectrochemistry*, 55, 161–164.
- Sunkari, V. G., Aranovitch, B., Portwood, N., & Nikoshkov, A. (2011). Effects of a low-intensity electromagnetic field on fibroblast migration and proliferation. *Electromagnetic Biology and Medicine*, 30, 80–85. doi:10.3109/15368378.2011.566774
- Suresh, S. J., Satish, A. V., & Choudhary, A. (2006). Influence of electric field on the hydrogen bond network of water. *The Journal of Chemical Physics*, 124, 74506. doi:10.1063/1.2162888
- Torgomyan, H., Hovnanyan, K., & Trchounian, A. (2013). *Escherichia coli* growth changes by the mediated effects after low-intensity electromagnetic irradiation of extremely high frequencies. *Cell Biochemistry and Biophysics*, 65, 445–454. doi:10.1007/s12013-012-9448-9
- Torgomyan, H., Kalantaryan, V., & Trchounian, A. (2011). Low intensity electromagnetic irradiation with 70.6 and 73 GHz frequencies affects *Escherichia coli* growth and changes water properties. *Cell Biochemistry and Biophysics*, 60, 275–281. doi:10.1007/s12013-010-9150-8
- Wertheimer, N., & Leeper, E. D. (1979). Electrical wiring configurations and childhood cancer. *American Journal of Epidemiology*, 109, 273–284. doi:10.1093/oxfordjournals.aje.a112681
- West, R. W., Hinson, W. G., Lyle, D. B., & Swicord, M. L. (1994). Enhancement of anchorage-independent growth in jb6 cells exposed to 60 hertz magnetic-fields. *Bioelectrochemistry and Bioenergetics*, 34, 39–43. doi:10.1016/0302-4598(94)80007-3



© 2019 The Author(s). This open access article is distributed under a Creative Commons Attribution (CC-BY) 4.0 license.

You are free to:

Share — copy and redistribute the material in any medium or format.

Adapt — remix, transform, and build upon the material for any purpose, even commercially.

The licensor cannot revoke these freedoms as long as you follow the license terms.

Under the following terms:

Attribution — You must give appropriate credit, provide a link to the license, and indicate if changes were made.

You may do so in any reasonable manner, but not in any way that suggests the licensor endorses you or your use.

No additional restrictions

You may not apply legal terms or technological measures that legally restrict others from doing anything the license permits.



***Cogent Biology* (ISSN: 2331-2025) is published by Cogent OA, part of Taylor & Francis Group.**

**Publishing with Cogent OA ensures:**

- Immediate, universal access to your article on publication
- High visibility and discoverability via the Cogent OA website as well as Taylor & Francis Online
- Download and citation statistics for your article
- Rapid online publication
- Input from, and dialog with, expert editors and editorial boards
- Retention of full copyright of your article
- Guaranteed legacy preservation of your article
- Discounts and waivers for authors in developing regions

**Submit your manuscript to a Cogent OA journal at [www.CogentOA.com](http://www.CogentOA.com)**

